



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Microplastic may increase uptake and bioaccumulation of organic pollutants in aquatic crustaceans

Frydkjær, Camilla Krogh; Iversen, Niels; Roslev, Peter

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Frydkjær, C. K., Iversen, N., & Roslev, P. (2016). *Microplastic may increase uptake and bioaccumulation of organic pollutants in aquatic crustaceans*. Abstract from SETAC Europe 26th Annual Meeting, Nantes, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

MO078 Microplastic may increase uptake and bioaccumulation of organic pollutants in aquatic crustaceans C.K. Frydkjær¹N.

Iversen, Aalborg University / Department of Chemistry and Bioscience;
P. Roslev, Aalborg University / Biology and Environmental Science.

Microplastic is causing an increasing environmental problem in aquatic environments where it has been shown to enter into the food chain.

Maybe even more important, microplastic may act as a contaminant trap and carrier of organic pollutants that could enhance bioaccumulation of these compounds in aquatic food chains. The uptake of microplastic in *Moina salina* was investigated using different concentrations and size fractions of microplastic particles. The amount of plastic taken up by the animals was determined by microscopy and the sublethal effects of microplastic on *Moina salina* were determined using video tracking and enzymatic assays. *Moina salina* ingested microplastic particles at various rates in these experiments. Microplastic particles similar to the ones used for the feeding experiments were allowed to adsorb ¹⁴C-phenanthrene, before these were fed to *Moina salina*. The radioactivity of the animals and of the microplastic particles was measured by liquid scintillation counting. The data obtained from these experiments were used to determine to which extent microplastic might act as a carrier of organic pollutants in aquatic environments.